

An International journal in English published online quarterly with aim to publish peer reviewed review and research articles in rapidly developing field of pharma and life sciences

We publish Research / Review article in the following subjects

Life Sciences

Branches : Agricultural Science, Biochemistry, Biology, Bioinformatics, Botany, Cytology, Cell biology, Chemistry, Ecology, Endocrinology, Entomology, Environmental Sciences, Food science and Technology, Genetics, Genomics & Proteomics, Immunobiology, Molecular biology, Marine Science, Microbiology, Neurobiology, Pathology, Physics, Physiology, Psychology, Veterinary Science, Zoology

Pharmaceutical Sciences

Branches : Biotechnology, Clinical and Hospital pharmacy, Herbal technology, Industrial Pharmacy, Immunology, International Regulatory Affairs, Medicine, Neuroscience, Novel drug delivery system, Nanotechnology, Pharmaceutics, Pharmacology & Toxicology, Pharmacognosy & Phytochemistry, Pharmacy practice, Pharmaceutical Engineering, Pharmaceutical Management, Pharmaceutical Analysis, Pharmaceutical Chemistry.



Impact Factor *0.672



Minneapolis , United States of America



U.S.A



Oxford, United Kingdom



Friedrich-Alexander-Universität Erlangen-Nürnberg



Erlangen, Germany



University of Reading

United Kingdom



ULRICHSWEB™
GLOBAL SERIALS DIRECTORY
USA (Washington DC)



Spring Hill College
Mamie and John Burke memorial library
USA (Alabama)



USA (Alabama)



USA (connecticut)



United kingdom



Japan



Japan



USA (Massachusetts)



DOSHISHA UNIVERSITY

Japan



Naropa University
Allen Ginsberg Library
USA (Colorado)



New York, NY, United States



Jacksonville State University
Houston Cole Library
USA (Alabama)



Illinois state university
Milner Library
USA (ILLINOIS)



Huntington College

Huntington College
Houghton Memorial library
USA (Alabama)



IJLPR would take care in making your article published without delay with your cooperation. IJLPR hopes that Researchers, Research scholars, Academician, Industrialists, Consultancy etc. would make use of this journal publication for the development of science and technology.

*Kindly visit Instruction to authors available at www.ijlpr.com for submission of manuscript for publication.

Any feed back / query kindly email to

editorofijlpr@rediffmail.com

editorijlpr@yahoo.com

or you can call +91 9908947749 / **+91 9676175127 / 9676175127**



SECOND DERIVATIVE SPECTROPHOTOMETRIC METHOD FOR SIMULTANEOUS DETERMINATION OF DOXYLAMINE SUCCINATE AND PYRIDOXINE HCL IN PHARMACEUTICAL FORMULATIONS.

AMAR KUMAR KASTURI¹, KANAKA RAJU MEDICHERLA², RAGHUVeer YAGANTI² AND N. BALAKRISHNAN^{3*}

¹Department of Chemistry, Pacific University, Udaipur, Rajasthan-313003, India.

²Department of Human Genetics, Andhra University, Visakhapatnam, Andhra Pradesh-530 003, India.

³Principal cum Professor, S A Raja Pharmacy College, Vadakkangulam-627116, Tamilnadu India.

ABSTRACT

Estimation of drugs in pharmaceutical formulations by using UV-Visible spectrophotometric method is one of the most frequently used techniques in pharmaceutical analysis. Simple and reliable second derivative spectrophotometric method was developed and validated for the simultaneous estimation of Doxylamine Succinate and Pyridoxine HCL in bulk and pharmaceutical formulations. The quantitative determination of the drugs was carried out using the second derivative values measured at 239 nm and 290 nm for Doxylamine Succinate and Pyridoxine Hcl respectively. The method has been developed and validated for the assay of Doxylamine Succinate and Pyridoxine HCL using methanol as diluents. Calibration graphs drawn at their wavelengths of determination were linear in the concentration range of 1-8 $\mu\text{g mL}^{-1}$ for Doxylamine Succinate and 12.5 - 100 $\mu\text{g mL}^{-1}$ for Pyridoxine Hcl. At 232 nm Doxylamine Succinate showed zero absorbance but Pyridoxine HCL had considerable amount of absorbance. Similarly at 290 nm Pyridoxine HCL showed zero absorbance but Doxylamine Succinate had considerable amount of absorbance. The limit of Quantification (LOQ) of Doxylamine Succinate and Pyridoxine HCL was found to be 0.6 $\mu\text{g mL}^{-1}$ and 1 $\mu\text{g mL}^{-1}$ respectively and the limit of detection (LOD) was found to be 0.08 $\mu\text{g mL}^{-1}$ and 0.3 $\mu\text{g mL}^{-1}$ respectively. Comparison second derivative spectrum of Doxylamine Succinate and Pyridoxine HCL in standard and drug formulation solutions showed that the wavelength of maximum absorbance did not change. The low relative standard deviation values indicate good precision and high recovery values indicate accuracy of the proposed method. Developed second derivative spectrophotometric method was simple, accurate, precise, specific, sensitive and reproducible, which can be directly and easily applied to pharmaceutical dosage forms.

KEYWORDS: *Spectrophotometric method, Second derivative, Doxylamine Succinate, Pyridoxine HCL.*



N. BALAKRISHNAN*

Principal cum Professor, S A Raja Pharmacy College, Vadakkangulam-627116, Tamilnadu India

Received on: 07-11-2018

Revised and Accepted on: 18-02-19

DOI: <http://dx.doi.org/10.22376/ijpbs/lpr.2019.9.2.P10-15>

INTRODUCTION

Doxylamine Succinate Ethanamine, N,N-dimethyl-2-[1-phenyl-1-(2-pyridinyl)ethoxy]-butanedioate (Fig.1) Doxylamine is a first-generation antihistamine.¹ It can be used by itself as a short-term sedative and in combination with other drugs to provide night-time allergy and cold relief.²⁻³ Doxylamine is also used in combination with the analgesics paracetamol (acetaminophen) and

codeine as an analgesic/calmativ preparation and is prescribed in combination with vitamin B6 (pyridoxine) to prevent morning sickness in pregnant women.⁴⁻⁵ Pyridoxine HCL 4-pyridinedimethanol, 5-hydroxy-6-methyl hydrochloride (Fig. 2). It is a water soluble vitamin and involved principally in amino acid, carbohydrate and fat metabolism.⁶⁻⁷ It is also required for the formation of haemoglobin.




Fig. 1
Doxylamine Succinate.




Fig. 2
Pyridoxine Hcl.

According to the literature survey it was found that few analytical methods such as Visible, UV, polarographic analysis and HPLC methods were reported for Pyridoxine Hcl and Doxylamine Succinate, but not even a single method has been reported on Second derivative spectroscopic method.⁸⁻⁹ The objective of this study was to develop and validate a simple and specific Second derivative spectrophotometric method for the simultaneous determination of Doxylamine Succinate and Pyridoxine HCL in combined dosage form.¹⁰⁻¹¹ Derivative spectroscopy has been widely used as a tool for quantitative analysis.¹²⁻¹³ This technique offers various advantages over the conventional spectrophotometric methods, such as discrimination of the sharp spectral features over the large bands and the enhancement of the resolution of the overlapping spectra.¹⁴⁻¹⁵ This method exhibited a precise, accurate and cost effective assay for these drugs in mixture.

MATERIALS AND METHODS

Selection of Solvent

The derivative spectra[s] of Doxylamine Succinate and Pyridoxine HCL in different solvents like

water, acetonitrile, sodium hydroxide and hydrochloric acid did not show any favorable zero crossing points, but when dissolved in methanol the derivative spectra[s] of both drugs showed zero crossing points. Hence methanol was selected as the solvent for the method.

Selection of derivative method

Though both first and second derivative spectra[s] showed zero crossing points in methanol solvent and their absorbance were considerably better, but the second derivative method was selected because the spectral characteristics and resolution were good in the second derivative spectra.

Selection of Wavelengths (Zero crossing points)

The zero crossing points of Doxylamine Succinate were 222, 232 and 235 nm and for Pyridoxine Hcl were 219, 239 and 241 nm. Out of these wavelengths 232 nm for Doxylamine Succinate and 239 nm for Pyridoxine Hcl were selected as the zero crossing points for the method based on their linearity data. At 232 nm Doxylamine Succinate showed zero absorbance but Pyridoxine Hcl had considerable amount of absorbance. Similarly at 239 nm Pyridoxine Hcl showed zero absorbance

but Doxylamine Succinate had considerable amount of absorbance.

Preparation of Standard stock solutions

Standard Doxylamine Succinate stock solution was prepared by dissolving 10mg of drug in 100 mL of methanol to get a concentration of 100 µg mL⁻¹. Standard Pyridoxine Hcl stock solution was prepared by dissolving 125 mg of drug in 100 mL of methanol to get a concentration of 1250 µg mL⁻¹. The standard solutions were prepared by dilution of the stock solution with methanol.

Chemicals

0.1N Hydrochloric acid, 0.1N Sodium hydroxide, Water, Acetonitrile, Methanol, Doxylamine Succinate and Pyridoxine HCL.

Instrument

Shimadzu UV-Visible spectrophotometer (model UV-1800).

Validation Parameters:

Linearity

To construct Beer lambert's law plot for Doxylamine Succinate (1-8 mL) in different concentrations (12.5, 25, 37.5, 50, 62.5, 75, 87.5 and 100 µg/mL) and Pyridoxine Hcl (0.1 - 0.8 mL) in different concentrations (1, 2, 3, 4, 5, 6, 7 and 8 µg/mL), were prepared by serial dilutions with methanol from the individual stock solutions. Then absorbance of the solution was measured at 232 nm for Pyridoxine Hcl and 239 nm for Doxylamine Succinate. The standard overlain spectrum of Doxylamine Succinate and Pyridoxine Hcl was shown in Fig. 3. The linearity values were shown in Table 1.




Fig.3
Overlain second derivative spectra of standard drugs (Doxylamine Succinate 2 µg mL⁻¹ and Pyridoxine Hcl 25 µg mL⁻¹).

Table. 1
Linearity

Parameters	Doxylamine Succinate	Pyridoxine Hcl
Concentration (µg/ mL)	1-8	12.5-100
Slope	0.001	0.0001
Intercept	0.0001	0.0001
Correlation coefficient (r ²)	0.999	0.999

Recovery studies

The recovery studies were carried out at three different levels i.e. 80%, 100% and 120%. To ensure the reliability of the above method, recovery studies were carried out by mixing a known

quantity of standard drug with the pre analyzed sample formulation and the contents were reanalyzed by the proposed method. The percentage recovery values were shown in Table. 2.

Table. 2
Recovery studies

Drug	Amount added	Amount recovered	% Recovery
Pyridoxine Hcl	12.5	12.47	99.76
	10	9.95	99.50
	15	14.94	99.60
DoxylamineHcl	1	0.991	99.10
	0.8	0.793	99.12
	1.2	1.19	99.16

Precision

The precision of the method was established by carrying out the analysis of the analytes using the

proposed method. The low value of standard deviation showed that the methods were precise. The results were shown in Table. 3.

Table. 3
Precision studies

Drug	Concentration	Intraday concentration measured	% RSD	Interday Values	%RSD
Pyridoxine Hcl	25	25.06	0.641	25.08	0.962
Doxylamine Succinate	2	2.03	0.874	2.04	0.891

Specificity

Comparison second derivative spectrum of Doxylamine Succinate and Pyridoxine Hcl in standard and drug formulation solutions showed that the wavelength of maximum absorbance did not change. According to the results obtained by recovery study, the derivative spectrophotometric method is able to access the analyte in presence of excipients and hence it can be considered as specific method.

LOD and LOQ

The limit of Detection (LOD) and limit of Quantification (LOQ) of the developed method were determined by injecting progressively low concentrations of the standard solutions. The LOQ of Doxylamine Succinate and Pyridoxine Hcl was found to be $0.6 \mu\text{g mL}^{-1}$ and $1 \mu\text{g mL}^{-1}$ respectively and the LOD was found to be $0.08 \mu\text{g mL}^{-1}$ and $0.3 \mu\text{g mL}^{-1}$ respectively.

Ruggedness

The ruggedness test of analytical assay method is defined as the degree of reproducibility of assay results

Obtained by the successful applications of assay over different time, day and among multiple analysts. The results showed no statistical differences suggesting that the developed method was rugged.

Preparation of Test Solution and estimation of Pyridoxine HCl and Doxylamine Succinate in formulation

For analysis of commercial formulations, 20 capsules were weighed, and powdered and weight equivalent to 12.5 mg of Pyridoxine HCl and 1 mg of Doxylamine Succinate was taken and transferred into a volumetric flask and made up to 100 mL with methanol, sonicated for 5 minutes, filtered and further diluted with methanol to get the concentration within the linearity range of respective drugs and measured the absorbance at 232 nm for Pyridoxine HCl and 239 nm for Doxylamine Succinate respectively. The formulation spectrum shown in Fig. 4. Then the amount of drug present in the formulation was calculated and results were shown in Table 4.

Table. 4
Analysis of formulation

Drug	Amount labeled (mg/mL)	Amount claimed (mg/mL)	% Label Claim	% Deviation
Pyridoxine Hcl	5 mg	4.95	99.00	(-) 1.0
Doxylamine Succinate	0.4 mg	0.397	99.25	(-) 0.75




Fig. 4

Second derivative spectra of formulation (Doxylamine Succinate 2 µg mL and Pyridoxine Hcl 25 µg mL)

RESULTS AND DISCUSSION

From the optical characteristics obtained with the proposed method it was found that the drug obeys linearity with in concentration range of 12.5 - 100 µg mL for Pyridoxine Hcl and 1-10 µg / mL for Doxylamine Succinate. From the precision studies, it was found that the percent relative standard deviation (% RSD) is less than 2%, which indicates that the method has good reproducibility. From the results of recovery studies, it was found that the percent recovery values of pure drug from the pre-analyzed solutions of formulations were in between 98.0-99.8%, which indicates the method is accurate and reveals that commonly used excipients and additives present in the pharmaceutical formulations did not interfere in the proposed method. The proposed method was simple, sensitive and reliable with good precision and accuracy. Hence, this method can be used for the routine analysis of Pyridoxine Hcl and Doxylamine Succinate in bulk samples and pharmaceutical formulations. Results and discussion part should be elaborated. Graph and tables, Please ensure that it should also come in results and discussion, Please rewrite this part

REFERENCES

1. SmitaNayak C, PreetiKulakarni V, Bhaskar V C V. Development and Validation of UV Spectrophotometric Method for Simultaneous Estimation of Doxylamine Succinate and

CONCLUSION

A convenient and rapid UV method has been developed for simultaneous estimation of Pyridoxine Hcl and Doxylamine Succinate in available dosage form. The assay provides a linear response across a wide range of concentrations. Low intra-day and interday % RSD coupled with excellent recoveries. Hence, this method can be easily and conveniently adopted for routine analysis of Pyridoxine Hcl and Doxylamine Succinate in pure form and its dosage forms.

AUTHOR'S CONTRIBUTION STATEMENT

Conceived and designed the experiments: AKK, NB. Performed the experiments: AKK, KM. Analyzed the data: AKK, NB. Contributed reagents/materials/analysis tools: NB, RY. Wrote the paper: AKK, KM.

CONFLICT OF INTEREST

Conflict of interest declared none

Pyridoxine Hydrochloride in Bulk and Tablet Dosage Form. Int J Pharm Pharm Sci. 2013;5(3):390-3.

Available

from:

- <http://www.ijppsjournal.com/Vol5Suppl3/7343.pdf>
2. Rajani SV, SowkarBaig I, Amulya R NP. A Validated Simultaneous Estimation of Doxylamine Succinate and Pyridoxine Hydrochloride by UV – Spectrophotometric Method in Bulk and Formulation. *Int J Pharm Res Anal.* 2014;4(2):139–43. Available from: <https://pdfs.semanticscholar.org/5831/cabe82438ad2cfddeb1663833a7d4937d25c.pdf>
 3. Gadsby R, Barnie-Adshead AM JC. Prospective study of nausea and vomiting during pregnancy. *Br J Gen Pr.* 1993;43:245–8. Available from: <http://bjgp.org/content/43/371/245.short>
 4. DV. F. Nausea and vomiting in pregnancy. *Am J Obs Gynecol.* 1968;102:135–75. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/4877794>
 5. Pope E, Maltepe C, Koren G. Comparing pyridoxine and doxylamine succinate-pyridoxine HCl for nausea and vomiting of pregnancy: A matched, controlled cohort study. *J Clin Pharmacol.* 2015;55(7):809–14. DOI; 10.1002/jcph.480
 6. Katsarov P, Gergov G, Alin A, Pilicheva B, Al-Degs Y, Simeonov V, et al. Advanced spectrophotometric chemometric methods for resolving the binary mixture of doxylamine succinate and pyridoxine hydrochloride. *Acta Pharm.* 2018;68(1):61–73. DOI: 10.2478/acph-2018-0008
 7. ACOG Issues Guidance on Treatment of Morning Sickness during Pregnancy. ACOG news release.; 2004. Available from: <https://m.acog.org/Patients/FAQs/Morning-Sickness-Nausea-and-Vomiting-of-Pregnancy>
 8. M.J.O’Neil. *The Merck Index: An Encyclopedia of Chemicals, Drugs, and Biologicals.* 14th ed. USA: Merck & Co; 2006. 2564 p. A Available from: <https://books.google.co.in/books?id=-AB8AQAACAAJ&dq=The+Merck+Index+14th+ed&hl=en&sa=X&ved=0ahUKEwj-3KrLk73gAhVDvo8KHcuuBt0Q6AEIKjAA>
 9. Ministry of Health and Family Welfare, Government of India. *The Indian Pharmacopoeia Commission.* 6th ed. Ghaziabad. New Delhi: Indian Pharmacopoeia; 2010. 2005 p. Available from: <https://www.scribd.com/document/279706354/Indian-Pharmacopoeia-2010>
 10. Rockville. *US Pharmacopoeia 27 (USP XXVII) National Formulary 22.* the United States Pharmacopoeial Q Convention; 2003. 672 p.
 11. Nataraj KS, Suvarna Y VG. Development and validation of method for simultaneous estimation of pyridoxine hydrochloride and doxylamine succinate in tablet dosage form by first order derivative spectroscopy. *Int J Pharm Pharm Sci.* 2012;5(1):388–90. Available from: <http://www.ijppsjournal.com/Vol5Issue1/6289.pdf>
 12. Argekar AP, Sawant JG. Simultaneous Determination of Pyridoxine Hydrochloride and Doxylamine Succinate from Tablets by Ion Pair Reversed-Phase High-Performance Liquid Chromatography (RP-HPLC). *Drug Dev Ind Pharm.* 1999;25(8):945–50. DOI: 10.1081/ddc-100102255
 13. Pathak A RS. Simultaneous determination of ternary mixture of doxylamine succinate, pyridoxine hydrochloride, and folic acid by the ratio spectra-zero-crossing, double divisor-ratio spectra derivative, and column high-performance liquid chromatographic methods. *J AOAC Int.* 2008;91(5):1059–69. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/18980119>
 14. Raza A, Ansari T, Niazi S RA. Spectrophotometric Determination of pyridoxine Hydrochloride (Vitamin B6) in Bulk and Tablets by Charge- Transfer Complexation with Chloranil. *J Chem Soc Pakistan.* 2007;29(1):33–6. Available from: https://www.researchgate.net/institution/National_Engineering_Services_Pakistan_Pvt_Limited/publications?nav=overview&ev=dep_publication
 15. International Conference of Harmonization ICH Harmonized Tripartite Guideline. In: *SpringerReference.* Springer-Verlag; DOI: 10.1007/springerreference_83218